

Accelerator Department
BROOKHAVEN NATIONAL LABORATORY
Associated Universities, Inc.
Upton, New York

AGS DIVISION TECHNICAL NOTE

No. 122

DECAY OF FLUCTUATION STATES IN STOCHASTIC COOLING

L.W. Smith
July 12, 1976

I. Introduction

Stochastic cooling¹ operates by sensing (and subsequently partially correcting) a statistical fluctuation in a finite sample of particles. I point out in this note that the mean life of such a statistical fluctuation is so short as to make the physical realization of stochastic cooling impractical.

II. Background

Suppose the beam sample, N , be divided into two parts with N_1 and N_2 particles on the positive and negative side, respectively, of the mean beam position. The sensed average position is

$$\bar{y} = \frac{N_1 - N_2}{N_1 + N_2} \cong \frac{N_1 - N_2}{N} s = \frac{n}{N} s$$

where s is a measure of the total beam size and we set $n \ll N$ so that we can take the denominator, N , to be a (quasi) constant.

If N_1 and N_2 are normally distributed, independent variables $N_1 - N_2$ is also normally distributed with mean zero and variance N . The probability density distribution function of n is

$$p(n) = \frac{1}{(2\pi N)^{\frac{1}{2}}} \exp\left(-\frac{n^2}{2N}\right),$$

and transforming to the variable, \bar{y} ,




Fig. 2 Bias Supply in NTM package




Fig. 3. Bias Supply Circuit Board

$$p(\bar{y}) = \frac{1}{s} \left(\frac{N}{2\pi} \right)^{\frac{1}{2}} \exp \left(-\frac{\bar{y}^2 N}{2s} \right)$$

with

$$\langle \bar{y}^2 \rangle = s^2 / N ,$$

i.e.

$$s^2 = \sigma^2 \cong \frac{\sum_i y_i^2}{N_T} \cong \frac{\sum_i y_i^2}{N} .$$

N_T is the total universe out of which N is a sample ($N_T > N$).

In a real beam we do not have a static situation such as drawing balls from an urn and the statistical fluctuation, $N \neq \langle N \rangle$, which produces the detected \bar{y} will persist for only some period of time, i.e. there are probability after effects.² The mean life of the fluctuation, \bar{y} , is the same as the mean life of the fluctuation, N .³

III. Calculation of Mean Life

We seek the probability that a particle will escape from the sample, N , which is within the azimuthal length from $-\frac{L}{2}$ to $+\frac{L}{2}$. In one revolution a particle will lag or lead a particle with central momentum p by

$$\ell = \eta C \frac{\Delta p}{p}$$

if its momentum is different from the central momentum by Δp ($\eta = \gamma_T^{-2} - \gamma^{-2}$).

We assume the sample to be uniformly distributed from $-\frac{L}{2}$ to $+\frac{L}{2}$. The probability of loss (in the positive ℓ direction) is equal to the probability of being in δx times the probability that $\ell > \frac{L}{2} - x$ integrated over the range of x which can contribute. The probability that $\ell > \frac{L}{2} - x$ is

$$\int_{L/2-x}^{\ell_m/2} d\ell p(\ell) \quad . \quad (1)$$

Where $p(\ell)$ is the probability density distribution of ℓ and $\ell_m/2$ is the maxi-

maximum (positive) value of ℓ . The probability of loss is then

$$P = \int_{x_0}^{L/2} \frac{dx}{L} \int_{\frac{\ell_m}{2}}^{\frac{\ell_m}{2} + x} d\ell p(\ell).$$

Where $x_0 = -L/2$ for $\ell_m/L \geq 2$ and $x_0 = \frac{1}{2}(L-\ell_m)$ for $\ell_m/L \leq 2$, i.e. we include in the integral only those particles which can be lost. If the distribution $p(\ell) = 1/\ell_m$ (uniform) Eq. 1 is equal to

$$\frac{1}{\ell_m} \left(\frac{\ell_m}{2} - \frac{L}{2} + x \right)$$

and we get (multiplying by 2 to include loss in the negative ℓ direction)

$$P = \begin{cases} \frac{\ell_m}{4L} & \text{for } \frac{\ell_m}{L} \leq 2 \\ 1 - \frac{L}{\ell_m} & \text{for } \frac{\ell_m}{L} \geq 2 \end{cases}$$

A physically more realistic $p(\ell)$ might be a parabolic one, i.e.

$$p(\ell) = \frac{3}{2} \frac{1}{\ell_m} - \frac{6}{\ell_m^3} \ell^2$$

and, as above, we get

$$P = \begin{cases} \frac{3}{16} \frac{\ell_m}{L} & \text{for } \frac{\ell_m}{L} \leq 2 \\ 1 - \frac{3}{2} \frac{L}{\ell_m} + \left(\frac{L}{\ell_m} \right)^3 & \text{for } \frac{\ell_m}{L} \geq 2 \end{cases}$$

and we see that P is not very sensitive to the exact form of $p(\ell)$.

It can be shown⁴ that the mean life of a state of fluctuation, N , for continuous observation is given by

$$T_N = 1/(N + \langle N \rangle) P_0$$

where $P(\delta t) = P_0 \delta t + O(\delta t^2)$. We see that the solutions of P for $\frac{\ell_m}{L} \leq 2$ satisfy this so we have ($N \cong \langle N \rangle$)

$$T_N \cong (\frac{1}{2} \langle N \rangle \frac{\ell_m}{L})^{-1}$$

Substituting $\langle N \rangle = N_T L/C$ and $\ell_m = 2\pi \frac{\Delta p_m}{p} C$ we get

$$T_N \cong (N_T \pi \frac{\Delta p_m}{p})^{-1} \text{ revolutions.}$$

For 10^{10} antiprotons stored in the ISA at 30 GeV with $\Delta p/p = \pm 10\%$ we get

$$T \cong 10^{-5} \text{ revolutions} \cong 10^{-10} \text{ sec}$$

or $\cong 3$ cm of azimuth.

There is another mechanism which contributes to the decay of the state of fluctuation, N . As above, assume the sampled beam to be divided into N_1 and N_2 particles on either side of the mean position. Then, due to betatron oscillation, in one revolution any particle will cross from region 1 to region 2 (or vice-versa) v times, and the mean life of the state N_1 is

$$\langle \langle N_1 \rangle \rangle \cong N_T / v$$

$$T_{N_1} \cong (2vN_1)^{-1} \cong T_{N_2}$$

so that the mean life of the state, N , using $\langle N_1 \rangle \cong \langle N_2 \rangle = 1/2 N$, is

$$T_N \cong (2vN)^{-1} \text{ revolutions.}$$

Substituting $N = 10^{10}$, $v \cong 25$ we get

$$T \cong 1/50 \times 10^{-5} \text{ revolutions.}$$

References

1. L.W. Smith, AGS Tech Note No. 120, and references therein.
2. S. Chandrasekhar, Stochastic Problems in Physics and Astronomy, Rev. Mod. Phys. 15, 1 (1943).
3. The mean life T_N is given by

$$\frac{1}{T_N} = \frac{1}{T_{N_1}} + \frac{1}{T_{N_2}}$$

or with m smaller slices

$$\frac{1}{T_N} = \sum_{i=1}^m \frac{1}{T_i}$$

4. Reference 2, p.53.

LWS/1sk

Distr: Dept. Admin.
AGS Division S&P
Scientific Staff in EP&S, ISA, and PD